


Chemists develop technique for extending nitrene reactions to three days

June 20 2024, by Bob Yirka



Top: orbital occupations and bond lengths of phenylnitrene (3). Bottom: Superimposed optimized geometries of $^3M^S$ FluidN (2) with the inset providing a close view of the central nitrene unit. Credit: *Science* (2024). DOI: 10.1126/science.adp4963

A team of chemists at the University of Bremen, in Germany has developed a new type of nitrene capable of slow reactions that can last for up to three days. Their [paper](#) is published in the journal *Science*.

Nitrenes (reactive intermediates and analogs of carbenes) are types of electrophile molecules that have a neutral atom bonded to a single other substituent. Because they have just six [electrons](#) in their valence level, they have brief reaction times, generally measured in the nanoseconds. That has made it difficult for chemists to use them in commercial applications.

In this new study, the researchers have found a way to dramatically slow nitrene reactions, by synthesizing a slow-reacting nitrene, possibly allowing for a whole new class of nitrenes that could be used in a wide variety of applications.

To achieve this feat, the research team employed the use of a M^S Fluid—a type of chemical scaffolding first built back in 2011 by another team. It has been used in other recent research efforts to stabilize other molecules, some of which are similar to nitrenes.

Using the scaffolding, the researchers were able to separate the

components in a reaction from accessing the [nitrogen atom](#), allowing for a much slower process. More specifically, they shone an [ultraviolet light](#) on a sample of $M^S\text{FluindN}_3$ (an azide precursor), which reduced it to $M^S\text{FluindN}$.

The researchers then subjected the newly synthesized nitrene to X-ray crystallography to learn more about its characteristics, followed by electron paramagnetic resonance spectroscopy and then superconducting quantum interference device magnetometry.

They confirmed that it had kept its spin-triplet ground state and persisted for up to three days. They also note that such imaging techniques are not normally possible with nitrene due to their fast reaction times.

The research team suggests their newly developed technique should allow for the synthesis of new transition metal complexes.

More information: Marvin Janssen et al, Synthesis of a stable crystalline nitrene, *Science* (2024). [DOI: 10.1126/science.adp4963](https://doi.org/10.1126/science.adp4963)

© 2024 Science X Network

Citation: Chemists develop technique for extending nitrene reactions to three days (2024, June 20) retrieved 21 June 2024 from <https://phys.org/news/2024-06-chemists-technique-nitrene-reactions-days.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.